

## LA-UR-16-27659

Approved for public release; distribution is unlimited.

Title: Semi-classical Electrodynamics: A Short Note

Author(s): Lestone, John Paul

Intended for: For general distribution to interested parties

Issued: 2016-10-05

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

## Semi-classical Electrodynamics: A Short Note

J. P. Lestone

Computational Physics Division, Los Alamos National Laboratory

Los Alamos, NM 87545, USA

September 27<sup>th</sup>, 2016

I have previously claimed the key to understanding the numerical value of the fine structure constant is near-field corrections which terminate integrals at low virtual photon energies, thus obverting an infrared divergence common to many QED calculations. I have since switched to a physics-based calculation of the near-field corrections, instead of the previously used educated guess. The relevant equations are presented here.

### Low-order Calculations

A previously reported physics-based calculation of the fine structure constant [1] has been refined. Instead of invoking black-hole physics, my reasoning has been mapped into a more QED-like picture, where virtual vacuum photons are assumed to isotropically scatter from isolated electrons with a cross section of  $\pi\lambda^2$ . In the presence of isolated electrons this scattering process violates conservation of energy and momentum, but is allowed for times given by the time-energy uncertainty principle. In the presence of a pair of electrons, the standard far-field vacuum photons are scattered, and then exchanged and rattle between the pair. The energy in the near field generates a repulsive inverse-square force and defines a numerical value for the fine structure constant  $\alpha \sim 1/137$ .

As previously reported, the fine structure constant can be expressed as

$$\alpha = \frac{1}{2\pi} \int_0^\infty \frac{f_{\text{nf}}^2(\varepsilon)}{\varepsilon(\exp(\varepsilon)-1)} d\varepsilon, \quad (1)$$

where  $f_{\text{nf}}(\varepsilon)$  is the near-field correction factor that scales the isolated scattering cross section to take into account near-field effects associated with the presence of a partner. It is squared because the buildup of the field energy between the pair is controlled by the scattering of far-field virtual photons into the region between the pair, and the scattering of the exchanging photons back into the far field. In the present short note I switch from a previous educated guess of the functional form of the near-field corrections, to a stronger explanation based on quantum mechanical reasoning associated with the overlap of the 3D wave functions of the scattered virtual photons surrounding each electron. The obtained low-order near-field correction is

$$f_{\text{nf}} = 1 - \exp^2\left(\frac{-\varepsilon^2\pi^2}{2^7}\right). \quad (2)$$

A document showing the derivation needs to wait until my return from an upcoming business trip.

Within my new picture, the anomalous magnetic moment of the electron can be estimated via

$$\frac{g-2}{2} = \frac{1}{4\pi} \int_0^\infty \frac{\exp(-\varepsilon)f_{\text{nf}}^2(\varepsilon)}{\varepsilon^3} d\varepsilon. \quad (3)$$

Using Eqs (1-3) the corresponding calculated quantities are  $\alpha=1/142.08$  and  $(g-2)/2=0.0011423$ .

### Possible Higher-order Corrections

Higher-order corrections to the proposed near-field based formulation of some QED processes are complex. Arguments exist (to be documented later) that suggest the near-field corrections to higher order might be of the form

$$f_{\text{nf}}^* = f_{\text{nf}} + (f_{\text{nf}})^n \exp^2\left(\frac{-\varepsilon^2\pi^2}{2^7}\right)/2, \quad (4)$$

where  $n$  is an unknown value of the order of unity. An estimate of  $n$  can be obtained by setting the relationship of our calculated  $\alpha$  and  $(g-2)/2$  to that known from QED [2],

$$\frac{g-2}{2} = \frac{\alpha}{2\pi} - 0.328478965579... \left(\frac{\alpha}{\pi}\right)^2 + 1.1812456587... \left(\frac{\alpha}{\pi}\right)^3 - \dots \quad (5)$$

The corresponding estimate is  $n=4.750479$ . This is only  $\sim 1$  part in  $10^4$  away from  $19/4$ . We do not know if any significance should be placed on the closeness of  $n$  to  $19/4$ . The corresponding predictions are  $(g-2)/2=0.001159478$  and  $\alpha=1/137.05664$ . These both differ from the known experimental values by  $\sim 1$  part in 7000, or  $\sim 3\alpha^2$  (relative). These differences are consistent with the accuracy expected of 4<sup>th</sup> order calculations with a missing 4<sup>th</sup> order term. If both Eqs (1) and (3) should be modified by the same 4<sup>th</sup> order correction scaling factor then this factor can be inferred via  $0.001159478 \div 0.001159652 = 0.9998498$ . The corresponding inferred  $\alpha$  is  $1/137.03605$ . This differs from the known value by  $\sim 1$  part in  $3 \times 10^6$  with a relative difference very close to  $\alpha^3$ . This last result may be fortuitous.

### Conclusions

These results support earlier suggestions that near-field corrections are the key to understanding the numerical value of the fine structure constant. More detailed studies of near-field effects should be pursued.

- [1] J. P. Lestone, Los Alamos National Laboratory Report, LA-UR-16-20131 (2016).
- [2] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, Phys. Rev. D **85**, 033007 (2012).